

May 2009

Gravitino Dark Matter and Non-Gaussianity

Tomo Takahashi¹, Masahide Yamaguchi²,
Jun'ichi Yokoyama^{3,4} and Shuichiro Yokoyama⁵

¹ *Department of Physics, Saga University, Saga 840-8502, Japan*

² *Department of Physics and Mathematics, Aoyama Gakuin University, Sagamihara
229-8558, Japan*

³ *Research Center for the Early Universe (RESCEU), Graduate School of Science, The
University of Tokyo, Tokyo 113-0033, Japan*

⁴ *Institute for the Physics and Mathematics of the Universe (IPMU), The University of
Tokyo, Kashiwa, Chiba, 277-8568, Japan*

⁵ *Department of Physics and Astrophysics, Nagoya University, Aichi 464-8602, Japan*

Abstract

We investigate density fluctuations in a scenario with gravitino dark matter in the framework of modulated reheating, which is known to generate large non-Gaussianity. We show that gravitino dark matter is disfavored in this framework. We also briefly discuss the case with the curvaton mechanism and some other possible dark matter scenarios.

Recent progress in high-precision observational cosmology with cosmic microwave background (CMB) radiation has motivated us to study the tiny nonlinear effects in the very early stage of the Universe when the amplitude of fluctuations was much smaller than unity. Among various observables to quantify non-Gaussianity of primordial fluctuations, the so-called non-linearity parameter f_{NL} , which is a stunted parameter to quantify the degrees of freedom of actual non-Gaussianity, has been widely discussed recently because observations of CMB is becoming to be able to measure its possible deviation from zero. From the recent result from WMAP5, the so-called local-type non-linearity parameter f_{NL} is constrained as $-9 < f_{\text{NL}} < 111$ [1] and $-4 < f_{\text{NL}} < 80$ [2] at 95 % confidence level (CL). Although purely Gaussian fluctuation with $f_{\text{NL}} = 0$ is perfectly consistent with the data, its observed central value is so much deviated from zero that future higher-precision observations may well confirm this local-type non-Gaussianity. Then the simplest models of single-field inflation [3] would be ruled out as a generation mechanism of primordial density/curvature fluctuations [4].

We can think of two mechanisms as an alternative to the inflaton to generate primordial fluctuations, modulated reheating [5] and the curvaton [6, 7], which are motivated in particle physics because there are many light fields in generic supersymmetric theories. Neither of them is necessary as long as the inflaton can be responsible for the generation of primordial fluctuations. But if the presence of large non-Gaussianity was established, they would be a target of serious study^{#1}. In fact, non-Gaussianity in these alternative scenarios has been studied already in [9, 10] for modulated reheating and in [11, 12] for the curvaton, and it has been shown that in both scenarios large local-type non-Gaussianity is possible depending on the values of model parameters and/or initial conditions.

The purpose of this Letter is to argue that confirmation of non-Gaussianity of local-type would not only rule out the simplest single-field slow-roll inflation models but also constrain the identity of the dark matter. That is, we show that the gravitino dark matter would confront difficulty in the two representative generation mechanisms of fluctuations mentioned above. Before proceeding to this main topic, for completeness, let us mention the relevance of the other type of non-Gaussianity, namely the equilateral-type, to the nature of the dark matter. It is generated due to nonlinear interactions during inflation. Typical models generating such non-Gaussianity are those making use of scalar fields with non-canonical kinetic terms [13]. They do not give us any information on the nature of the dark matter, because the inflaton is responsible for the fluctuation in this case. Observationally, the non-linearity parameter of the equilateral-type, $f_{\text{NL}}^{\text{equil}}$, has been much less constrained than the local-type, and as long as the former is concerned, there is no hint of non-Gaussianity so far. We therefore concentrate on the local-type non-Gaussianity which we denote simply by f_{NL} as above.

In models with a supersymmetric extension of the standard model, the gravitino, the spin-3/2 superpartner of the graviton, arises when the global supersymmetry is promoted to a local symmetry. If it is the lightest supersymmetric particle (LSP) and stable be-

^{#1} Other phenomenological generation mechanisms of large local-type non-Gaussianity include Refs. [8].

cause of the R -parity conservation, it can be a good candidate of dark matter (DM) of the Universe. Gravitinos can be produced in scattering processes of particles in the thermal bath and its thermal relic density depends on the efficiency of the production, which is sensitive to the reheating temperature T_R after inflation. Its relic abundance is given as $Y_{3/2} \equiv n_{3/2}/s \propto T_R$ where $Y_{3/2}$ is the yield from the thermal production, $n_{3/2}$ and s are the number density of gravitinos and the entropy of the universe [14]. Recently, it was pointed out that gravitinos are also produced non-thermally from the decay of heavy scalar fields like an inflaton or moduli if they acquire non-vanishing expectation values [15]. The abundance of gravitinos produced non-thermally also depends on the reheating temperature, in fact, is proportional to the inverse of the reheating temperature. Then, when one considers the modulated reheating scenario as a generation mechanism of primordial fluctuations, the dependence on T_R would result in the generation of isocurvature fluctuations in gravitino DM since the modulated reheating scenario indicates that $\delta T_R/T_R \neq 0$. Such isocurvature fluctuations can be strongly constrained from cosmological observations such as CMB. From WMAP5, the ratio of isocurvature fluctuations to curvature fluctuations, which is denoted by α , is constrained as $\alpha < 0.072$ (axion type; uncorrelated) and $\alpha < 0.0041$ (curvaton type; anti-correlated) for WMAP+BAO+SN at the 95% CL [1] (see also Ref. [16] for the constraint on isocurvature fluctuations from the non-Gaussianity).

One may classify the origin of gravitino DM to the following two cases in the context of the modulated or the curvaton scenario. The first one is that gravitino can be produced at the decay of the inflaton through the rescattering process of thermal plasma (thermal production) or directly by the inflaton decay (non-thermal production). In this case, such gravitino DM results in the generation of large isocurvature fluctuations not only in the modulated scenario but also in the curvaton scenario [11]. The other one is the case that the curvaton mechanism is operative and gravitino DM is mostly produced from the decay of the curvaton. In this case, isocurvature fluctuations are suppressed if and only if the curvaton practically dominates the energy density of the Universe just before its decay [11]. However, such a case cannot generate the large non-Gaussianity. Thus when large non-Gaussianity is confirmed in the future, a scenario with gravitino DM may not be a viable candidate as the main constituent of the DM in the Universe.

In this Letter, we discuss in detail the viability of the gravitino DM scenario when non-Gaussianity is large, particularly focusing on the modulated reheating scenario [5] since the issues of isocurvature fluctuations in this scenario is relatively unexplored. As shown below, the gravitino DM acquires large isocurvature fluctuations in this scenario, which is too large to be consistent with current observations of CMB.

Now we briefly review the non-linearity parameter which characterizes non-Gaussianity of the fluctuations. For this purpose, it is fashionable to write the comoving curvature perturbation, ζ , up to the second order to discuss the bispectrum as

$$\zeta = \zeta_{(1)} + \frac{3}{5}f_{\text{NL}}\zeta_{(1)}^2, \quad (1)$$

where $\zeta_{(1)}$ is the Gaussian part of curvature perturbation and f_{NL} characterizes the size of non-Gaussianity observable in the bispectrum. With this parameterization, the definition

of the bispectrum B_ζ is given by

$$\langle \zeta_{\vec{k}_1} \zeta_{\vec{k}_2} \zeta_{\vec{k}_3} \rangle = (2\pi)^3 B_\zeta(k_1, k_2, k_3) \delta(\vec{k}_1 + \vec{k}_2 + \vec{k}_3). \quad (2)$$

The leading contribution of the bispectrum B_ζ can be written as

$$B_\zeta(k_1, k_2, k_3) = \frac{6}{5} f_{\text{NL}} \left(P_\zeta(k_1) P_\zeta(k_2) + P_\zeta(k_2) P_\zeta(k_3) + P_\zeta(k_3) P_\zeta(k_1) \right). \quad (3)$$

Here P_ζ is the power spectrum which is defined as

$$\langle \zeta_{\vec{k}_1} \zeta_{\vec{k}_2} \rangle = (2\pi)^3 P_\zeta(k_1) \delta(\vec{k}_1 + \vec{k}_2). \quad (4)$$

In order to evaluate the curvature perturbations ζ , we adopt the δN formalism [17]. The curvature perturbation on sufficiently large scales ζ at time t_f is equal to the perturbation in time integral of the local expansion from an initial flat hypersurface ($t = t_i$) to the final uniform energy density hypersurface. Since the local expansion on sufficiently large scales can be approximated by the expansion of the unperturbed Friedmann universe, the curvature perturbation ζ is evaluated as

$$\zeta(t_f, \vec{x}) = N(t_i, t_f, \vec{x}) - (\text{spatial average}), \quad (5)$$

where the number of e -folds $N(t_i, t_f, \vec{x})$ is defined by the time integral of the local Hubble parameter as,

$$N(t_i, t_f, \vec{x}) = \int_{t_i}^{t_f} H(t, \vec{x}) dt. \quad (6)$$

In the δN formalism, taking $t_i = t_*$ which is a certain time well after the relevant scale crossed the horizon scale, the curvature fluctuations from fluctuations of scalar fields are given as,

$$\zeta = N_a \delta\varphi_*^a + \frac{1}{2} N_{ab} \delta\varphi_*^a \delta\varphi_*^b + \dots, \quad (7)$$

up to the second order, where $\delta\varphi_*^a$ is the long-wave frozen part of the fluctuation of the scalar field φ^a on the initial flat hypersurface at $t = t_*$ and N_a and N_{ab} are given by

$$N_a \equiv \frac{\partial N}{\partial \varphi^a}, \quad N_{ab} \equiv \frac{\partial^2 N}{\partial \varphi^a \partial \varphi^b}. \quad (8)$$

Here the summation is implied for the repeated indices.

The non-linearity parameter is then given by

$$\frac{6}{5} f_{\text{NL}} = \frac{N_{ab} N^a N^b}{(N_c N^c)^2}, \quad (9)$$

$$(10)$$

where the indices are lowered and raised by using the Kronecker's delta δ^{ab} .

Using δN formalism, we can evaluate the curvature perturbations in the modulated reheating scenario. In this scenario, fluctuations, $\delta\sigma$, of some modulus, σ , renders fluctuations in the decay rate of the inflaton. To evaluate ζ from the fluctuations of the modulus, we need to follow the background evolution from the epoch after the inflaton begins to oscillate around the minimum of the potential whose shape is assumed to be quadratic below. Then the energy density of the oscillating inflaton field behaves like matter, thus the background equations after inflation are governed by the following equations:

$$\frac{d\rho_\phi}{dN} + 3\rho_\phi = -\frac{\Gamma}{H}\rho_\phi, \quad (11)$$

$$\frac{d\rho_r}{dN} + 4\rho_r = \frac{\Gamma}{H}\rho_\phi, \quad (12)$$

$$H^2 = \frac{1}{3}(\rho_\phi + \rho_r), \quad (13)$$

where Γ is the σ -dependent decay rate of the inflaton and ρ_ϕ and ρ_r are energy densities of inflaton ϕ and radiation, respectively. We assume the energy density of the modulus is negligible throughout. The reduced Planck scale is set to be unity here.

By solving the above equations from the end of inflation to the completion of reheating with the initial condition $\rho_\phi(0) = \rho_0 = 3H_0^2$ and $\rho_r(0) = 0$, we can obtain a relation between N and Γ . The number of e -folds, N , elapsed while the Hubble parameter drops from H_0 to H_f ($H_f \ll \Gamma$) can be written formally as [10]

$$N = \frac{1}{2} \log \frac{H_0}{H_f} + Q \left(\frac{\Gamma}{H_0} \right), \quad (14)$$

where the function Q is defined by

$$\exp \left[4Q(\Gamma/H_0) \right] \equiv \int_0^\infty dN' \frac{\Gamma}{H(N')} e^{4N' \frac{\rho_\phi(N')}{\rho_0}}. \quad (15)$$

On dimensional grounds Q depends only on the combination of $x \equiv \Gamma/H_0$. Let us consider the typical case that the final stage of reheating is governed by a perturbative decay of the inflaton after a certain period of field-oscillation dominated stage. Then x is much smaller than unity and we find the following analytic form of $Q(x)$ [10],^{#2}

$$Q(x) = -\frac{1}{6} \log x + \mathcal{O}(x), \quad x \ll 1. \quad (16)$$

In the modulated reheating scenario, the curvature perturbation ζ is evaluated as [10]

$$\zeta(N_F) = Q_\sigma \delta\sigma_* + \frac{1}{2} Q_{\sigma\sigma} \delta\sigma_*^2, \quad (17)$$

^{#2} The behavior of the inflaton after inflation depends on the potential shape and the decay rate. Here we consider the case that the inflaton behaves like matter assuming the effective potential around the minimum is quadratic and oscillates enough before its decay. For other cases, some coefficients are changed but the essential results remain intact.

up to the second order with

$$Q_\sigma \equiv Q'(x) \frac{\Gamma'(\sigma)}{H_0} , \quad (18)$$

$$Q_{\sigma\sigma} \equiv Q'(x) \frac{\Gamma''(\sigma)}{H_0} + Q''(x) \left(\frac{\Gamma'(\sigma)}{H_0} \right)^2 , \quad (19)$$

where $Q'(x) = dQ(x)/dx$ and $\Gamma'(\sigma) = d\Gamma(\sigma)/d\sigma$.

From the above equations, we can obtain the following expression for the non-linearity parameter

$$\frac{6}{5} f_{\text{NL}} \simeq 6 \left(1 - \frac{(\Gamma''(\sigma)/\Gamma(\sigma))}{(\Gamma'(\sigma)/\Gamma(\sigma))^2} \right) \quad (20)$$

in the case $x \ll 1$.

Hence, if we adopt proper form of the decay rate, we can realize a large non-Gaussianity in the modulated reheating scenario. As an example, let us consider the case that the decay rate depends on σ as

$$\Gamma(\sigma) = \Gamma_0 \left[1 + A \frac{\sigma}{M} + B \left(\frac{\sigma}{M} \right)^2 \right] , \quad (21)$$

where A and B are some coefficients, M is a some energy scale and we have assumed $\sigma/M \ll 1$ and truncated at the second order in σ/M . In this model, we find an simple expression for f_{NL} as

$$\frac{6}{5} f_{\text{NL}} \simeq 6 \left(1 - \frac{2B}{A^2} \right) . \quad (22)$$

Thus, if we take $|B|/A^2 \gg 1$ and $B < 0$, we can obtain a large and positive value of f_{NL} .

Let us move on to the isocurvature fluctuations from gravitino DM in the modulated reheating scenario. As mentioned before, there are several distinct ways to produce gravitinos. One is the scattering of particles in the thermal plasma and their relic abundance is evaluated as [14]

$$Y_{3/2} \equiv \frac{n_{3/2}}{s} \simeq 10^{-12} \sum_i g_i^2 \left(1 + \frac{m_{Gi}^2}{3m_{3/2}^2} \right) \left(\frac{T_R}{10^{10} \text{ GeV}} \right) \quad (23)$$

in terms of the yield, where $n_{3/2}$ is the number density of gravitino and s is the entropy density. In the right hand side, T_R is the reheating temperature, $m_{3/2}$ is the gravitino mass, m_{Gi} is the gaugino masses for i -th generation, and g_i is the gauge coupling^{#3}.

^{#3} In fact, the gauge coupling and the gaugino masses mildly depend on the reheating temperature T_R . However we neglect the weak dependences in the following discussions.

Notice that the relic abundance is proportional to the reheating temperature T_R . In the modulated reheating scenario, the reheating temperature fluctuates in space, which implies that the number density of gravitinos per entropy also fluctuates. Then, if the gravitinos constitute DM of the Universe, isocurvature fluctuations are generated. In order to discuss it more quantitatively, we relate the fluctuations of the number density of gravitinos to the curvature fluctuations as

$$S_{3/2} \equiv \frac{\delta(n_{3/2}/s)}{n_{3/2}/s}. \quad (24)$$

In the modulated reheating scenario, $\delta T_R/T_R \neq 0$, which implies that a significant isocurvature fluctuation is generated as

$$S_{3/2} = \frac{\delta T_R}{T_R}. \quad (25)$$

Since T_R is proportional to $\Gamma^{1/2}$, we find $\delta T_R/T_R = \delta\Gamma/(2\Gamma) = \delta x/(2x)$. Then using the expression for the curvature perturbation (17), $S_{3/2}$ can be related to ζ as

$$S_{3/2} = \frac{\delta T_R}{T_R} = \frac{1}{2xQ'(x)}\zeta \simeq -3\zeta, \quad (26)$$

when $x \ll 1$. Thus we have $S_{3/2}/\zeta \simeq -3$, which is totally (anti)-correlated to the curvature perturbation and already contradicts with the current observation [1]. This shows that a scenario with thermally produced gravitino DM cannot be viable in the modulated reheating scenario, in which large non-Gaussianity can be generated, because of too large isocurvature fluctuations.

In fact, a similar argument also holds when gravitinos are produced non-thermally from the decay of some heavy scalar field such as the inflaton or moduli in the case they acquire a nonvanishing vacuum expectation value [15], which dominates over the other channels [18]. In this case, the yield can be written as

$$\frac{n_{3/2}}{s} = \frac{3}{2}B_{3/2}\frac{T_R}{m_{3/2}}, \quad (27)$$

where $B_{3/2}$ is the branching ratio of the decay into gravitinos. When gravitinos are produced from the jets, $B_{3/2}$ should be understood as those including its multiplicity. Since the reheating temperature is related to the decay rate of the inflaton as $T_R \propto \Gamma^{1/2}$, then $B_{3/2} \propto T_R^{-2}$, we have the T_R -dependence of the $n_{3/2}/s$, in the case of non-thermal production, as

$$\frac{n_{3/2}}{s} \propto \frac{1}{T_R}. \quad (28)$$

Thus isocurvature fluctuations are also generated when gravitinos are produced from some heavy scalar decay, which can be written as

$$S_{3/2} = -\frac{\delta T_R}{T_R}, \quad (29)$$

which is different in sign but the same in size as in the case that gravitinos are produced thermally. Note that this isocurvature fluctuation has totally *positive* correlation to the curvature perturbation. Hence if DM consists of gravitinos, large isocurvature fluctuations are generically produced in the modulated reheating scenario, which is inconsistent with cosmological observations such as CMB.

There is another well-known mechanism to generate large local-type non-Gaussianity, called the curvaton mechanism. In this scenario, gravitinos are produced from the decays of both the inflaton and the curvaton. As discussed in Ref. [11, 19], in the former case, gravitino DM is disfavored because of too large isocurvature fluctuations. On the other hand, in the latter case, gravitino DM is permitted only when the curvaton almost dominates the energy density of the universe to avoid large isocurvature fluctuations, in which the non-Gaussianity is small. Then, it is concluded that gravitino DM is disfavored also in the curvaton scenario if the large non-Gaussianity is observed ^{#4}. Therefore, gravitino dark matter is disfavored once large local-type non-Gaussianity of primordial fluctuations is observed.

Some comments are in order. First gravitinos can also be produced by the decay of the next-to-the-lightest supersymmetric particle (NLSP). However, the detailed calculations show that the constraints from BBN are severer than those from the overclosure of the universe, irrespective of the particle kind of NLSP as long as it is the MSSM particle [21]. Therefore, gravitino DM can be realized only for the case that most of gravitinos should be produced much earlier, either by the scattering processes during reheating or by the non-thermal decay of heavy scalars, both of which we paid attention to in this Letter.

Second, a similar argument also holds for the case where LSPs, which are DM, are produced from the decay of gravitinos. In this case, LSPs carry large isocurvature fluctuations originating from gravitinos, which again contradicts with the present constraint. Thus, DMs originating from gravitinos are also disfavored once large local-type non-Gaussianity is detected ^{#5}.

Finally we comment on the case of axino DM. Axinos are produced by thermal scattering during reheating and their abundance also depends on the reheating temperature [22–25]. Therefore, as with the case of gravitinos, if such axinos become DM, they also acquire large isocurvature fluctuations in the modulated reheating scenario, which again contradicts with the present observational constraints. Unlike the gravitino dark matter, however, non-thermally produced axinos by the decay of NLSP [24–26] or Q-balls [27] can be dominant with avoiding the BBN constraints. For such cases, isocurvature constraints can be waived.

Acknowledgments:

We thank Takahiro Tanaka for the collaboration at the early stage. We are also grateful to Takashi Hamazaki, Masahiro Kawasaki, Kazunori Kohri, and Fuminobu Takahashi for

^{#4} For the case of baryonic isocurvature fluctuations, see Ref. [20].

^{#5} We thank M. Kawasaki and F. Takahashi for pointing out this possibility.

useful discussions. This work is supported by JSPS Grant-in-Aid for Scientific research, No. 19740145 (T.T.), No. 19340054 (J.Y.), and No. 21740187 (M.Y.). S.Y. is supported in part by Grant-in-Aid for Scientific Research on Priority Areas No. 467 “Probing the Dark Energy through an Extremely Wide and Deep Survey with Subaru Telescope”. He also acknowledges the support from the Grand-in-Aid for the Global COE Program “Quest for Fundamental Principles in the Universe: from Particles to the Solar System and the Cosmos ” from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan. We would like to thank the organizers of the IPMU workshop on “Focus week on non-Gaussianities in the sky” and the GCOE/YITP workshop YITP-W-09-01 on “Non-linear cosmological perturbations” for their hospitality, during which a part of this work is done.

References

- [1] E. Komatsu *et al.* [WMAP Collaboration], arXiv:0803.0547 [astro-ph].
- [2] K. M. Smith, L. Senatore and M. Zaldarriaga, arXiv:0901.2572 [astro-ph].
- [3] A.H. Guth, Phys. Rev. **D23**, 347 (1981); K. Sato, Mon. Not. Roy. Astron. Soc. **195**, 467 (1981); A.A. Starobinsky Phys. Lett. **91B**, 99 (1980); A. D. Linde, Phys. Lett. **108B**, 389 (1982); A. Albrecht and P. J. Steinhardt, Phys. Rev. Lett. **48**, 1220 (1982); A.D. Linde, Phys. Lett. B **129**, 177 (1983).
- [4] S.W. Hawking, Phys. Lett. **115B**, 295 (1982); A.A. Starobinsky, Phys. Lett. **117B**, 175 (1982); A.H. Guth and S-Y. Pi, Phys. Rev. Lett. **49**, 1110 (1982).
- [5] G. Dvali, A. Gruzinov and M. Zaldarriaga, Phys. Rev. D **69**, 023505 (2004) [arXiv:astro-ph/0303591].
L. Kofman, arXiv:astro-ph/0303614.
- [6] S. Mollerach, Phys. Rev. D **42**, 313 (1990).
A. D. Linde and V. F. Mukhanov, Phys. Rev. D **56**, 535 (1997) [arXiv:astro-ph/9610219].
- [7] K. Enqvist and M. S. Sloth, Nucl. Phys. B **626**, 395 (2002) [arXiv:hep-ph/0109214];
D. H. Lyth and D. Wands, Phys. Lett. B **524**, 5 (2002) [arXiv:hep-ph/0110002];
T. Moroi and T. Takahashi, Phys. Lett. B **522**, 215 (2001) [Erratum-ibid. B **539**, 303 (2002)] [arXiv:hep-ph/0110096].
- [8] E. W. Kolb, A. Riotto and A. Vallinotto, Phys. Rev. D **73**, 023522 (2006) [arXiv:astro-ph/0511198].
D. H. Lyth, JCAP **0511**, 006 (2005) [arXiv:astro-ph/0510443].

- L. Alabidi and D. Lyth, JCAP **0608**, 006 (2006) [arXiv:astro-ph/0604569].
M. Sasaki, Prog. Theor. Phys. **120**, 159 (2008) [arXiv:0805.0974 [astro-ph]].
- [9] M. Zaldarriaga, Phys. Rev. D **69**, 043508 (2004) [arXiv:astro-ph/0306006].
A. Mazumdar, Phys. Rev. Lett. **92**, 241301 (2004) [arXiv:hep-ph/0306026].
K. Ichikawa, T. Suyama, T. Takahashi and M. Yamaguchi, Phys. Rev. D **78**, 063545 (2008) [arXiv:0807.3988 [astro-ph]].
- [10] T. Suyama and M. Yamaguchi, Phys. Rev. D **77**, 023505 (2008) [arXiv:0709.2545 [astro-ph]].
- [11] D. H. Lyth, C. Ungarelli and D. Wands, Phys. Rev. D **67**, 023503 (2003) [arXiv:astro-ph/0208055].
D. H. Lyth and D. Wands, Phys. Rev. D **68**, 103516 (2003) [arXiv:astro-ph/0306500].
- [12] N. Bartolo, S. Matarrese and A. Riotto, Phys. Rev. D **69**, 043503 (2004) [arXiv:hep-ph/0309033].
K. Enqvist and S. Nurmi, JCAP **0510**, 013 (2005) [arXiv:astro-ph/0508573].
K. A. Malik and D. H. Lyth, JCAP **0609**, 008 (2006) [arXiv:astro-ph/0604387].
M. Sasaki, J. Valiviita and D. Wands, Phys. Rev. D **74**, 103003 (2006) [arXiv:astro-ph/0607627].
Q. G. Huang, arXiv:0801.0467 [hep-th].
K. Ichikawa, T. Suyama, T. Takahashi and M. Yamaguchi, Phys. Rev. D **78**, 023513 (2008) [arXiv:0802.4138 [astro-ph]].
K. Enqvist and T. Takahashi, JCAP **0809**, 012 (2008) [arXiv:0807.3069 [astro-ph]].
Q. G. Huang, JCAP **0811**, 005 (2008) [arXiv:0808.1793 [hep-th]].
Q. G. Huang and Y. Wang, JCAP **0809**, 025 (2008) [arXiv:0808.1168 [hep-th]].
- [13] M. Alishahiha, E. Silverstein and D. Tong, Phys. Rev. D **70**, 123505 (2004) [arXiv:hep-th/0404084].
N. Arkani-Hamed, P. Creminelli, S. Mukohyama and M. Zaldarriaga, JCAP **0404**, 001 (2004) [arXiv:hep-th/0312100].
- [14] J. R. Ellis, J. E. Kim and D. V. Nanopoulos, Phys. Lett. B **145**, 181 (1984).
T. Moroi, H. Murayama and M. Yamaguchi, Phys. Lett. B **303**, 289 (1993).
M. Bolz, A. Brandenburg and W. Buchmuller, Nucl. Phys. B **606**, 518 (2001) [Erratum-ibid. B **790**, 336 (2008)] [arXiv:hep-ph/0012052].
J. Pradler and F. D. Steffen, Phys. Rev. D **75**, 023509 (2007) [arXiv:hep-ph/0608344].

- [15] S. Nakamura and M. Yamaguchi, Phys. Lett. B **638**, 389 (2006) [arXiv:hep-ph/0602081].
T. Asaka, S. Nakamura and M. Yamaguchi, Phys. Rev. D **74**, 023520 (2006) [arXiv:hep-ph/0604132].
M. Kawasaki, F. Takahashi and T. T. Yanagida, Phys. Lett. B **638**, 8 (2006) [arXiv:hep-ph/0603265]; Phys. Rev. D **74**, 043519 (2006) [arXiv:hep-ph/0605297].
M. Endo, K. Hamaguchi and F. Takahashi, Phys. Rev. Lett. **96**, 211301 (2006) [arXiv:hep-ph/0602061].
M. Endo, F. Takahashi and T. T. Yanagida, Phys. Rev. D **76**, 083509 (2007) [arXiv:0706.0986 [hep-ph]].
- [16] C. Hikage, K. Koyama, T. Matsubara, T. Takahashi and M. Yamaguchi, arXiv:0812.3500 [astro-ph].
- [17] A. A. Starobinsky, JETP Lett. **42** (1985) 152 [Pisma Zh. Eksp. Teor. Fiz. **42** (1985) 124].
Y. Nambu and A. Taruya, Class. Quant. Grav. **13**, 705 (1996) [arXiv:astro-ph/9411013].
M. Sasaki and E. D. Stewart, Prog. Theor. Phys. **95**, 71 (1996).
M. Sasaki and T. Tanaka, Prog. Theor. Phys. **99**, 763 (1998).
H. Kodama and T. Hamazaki, Phys. Rev. D **57**, 7177 (1998) [arXiv:gr-qc/9712045].
D. H. Lyth, K. A. Malik, and M. Sasaki, J. Cosmol. Astropart. Phys. **05**, 004 (2005).
T. Hamazaki, Phys. Rev. D **78**, 103513 (2008) [arXiv:0811.2366 [astro-ph]].
- [18] M. Hashimoto, K. I. Izawa, M. Yamaguchi and T. Yanagida, Prog. Theor. Phys. **100**, 395 (1998) [arXiv:hep-ph/9804411];
K. Kohri, M. Yamaguchi and J. Yokoyama, Phys. Rev. D **70**, 043522 (2004) [arXiv:hep-ph/0403043].
- [19] M. Lemoine, J. Martin and J. Yokoyama, arXiv:0903.5428 [astro-ph.CO]; arXiv:0904.0126 [astro-ph.CO].
- [20] T. Moroi and T. Takahashi, Phys. Lett. B **671**, 339 (2009) [arXiv:0810.0189 [hep-ph]].
- [21] M. Kawasaki, K. Kohri, T. Moroi and A. Yotsuyanagi, Phys. Rev. D **78**, 065011 (2008) [arXiv:0804.3745 [hep-ph]].
- [22] K. Rajagopal, M. S. Turner and F. Wilczek, Nucl. Phys. B **358**, 447 (1991).

- [23] L. Covi, L. Roszkowski and M. Small, JHEP **0207**, 023 (2002) [arXiv:hep-ph/0206119].
- [24] L. Covi, H. B. Kim, J. E. Kim and L. Roszkowski, JHEP **0105**, 033 (2001) [arXiv:hep-ph/0101009].
- [25] A. Brandenburg and F. D. Steffen, JCAP **0408**, 008 (2004) [arXiv:hep-ph/0405158].
A. Freitas, F. D. Steffen, N. Tajuddin and D. Wyler, arXiv:0904.3218 [hep-ph].
- [26] L. Covi, J. E. Kim and L. Roszkowski, Phys. Rev. Lett. **82**, 4180 (1999) [arXiv:hep-ph/9905212].
- [27] L. Roszkowski and O. Seto, Phys. Rev. Lett. **98**, 161304 (2007) [arXiv:hep-ph/0608013].
O. Seto and M. Yamaguchi, Phys. Rev. D **75**, 123506 (2007) [arXiv:0704.0510 [hep-ph]].